

# 49 DOSES OF 14 VACCINES BEFORE AGE 6? 69 DOSES OF 16 VACCINES BY AGE 18?

*Before you take the risk, find out what it is.*

<b>BIRTH (12 hours)</b>	<b>2 MONTHS</b>	<b>4 MONTHS</b>	<b>6 MONTHS</b>	<b>7 MONTHS</b>	<b>12 - 18 MONTHS</b>	<b>2 - 6 YEARS</b>	<b>7-18 YEARS</b>
Hepatitis B	Diphtheria	Diphtheria	Diphtheria	Influenza	Diphtheria	Diphtheria	Diphtheria
	Tetanus	Tetanus	Tetanus		Tetanus	Tetanus	Tetanus
	Pertussis	Pertussis	Pertussis		Pertussis	Pertussis	Pertussis
	Polio	Polio	Polio		Measles	Polio	Influenza (12)
	HIB	HIB	Rotavirus		Mumps	Measles	HPV (3)
	Rotavirus	Rotavirus	Hepatitis B		Rubella	Mumps	Meningococcal (2)
	Hepatitis B	PCV	PCV		HIB	Rubella	
	PCV		Influenza		PCV	Varicella	
					Varicella	Influenza (5)	
					Hepatitis A (2)		



**VACCINE INGREDIENTS:** Different vaccines contain different ingredients including lab altered live or inactivated viruses and bacteria, chemicals, metals, proteins, antibiotics and human, animal and insect DNA and RNA. Learn more at [NVIC.org](http://NVIC.org).

If You Vaccinate Your Child,  
**Learn How to Recognize the Signs  
 and Symptoms of Vaccine Reactions\***



VACCINE REACTIONS	MOTHER'S DESCRIPTIONS
<b>High Fever (over 103° F)</b>	"His temperature was 105 degrees. I had to put cool towels on him to bring the fever down."
<b>Skin (hives, rashes, swelling)</b>	"There was a big, hot swollen lump at the site of the shot that stayed for weeks."
<b>High Pitched Screaming</b>	"It was a pain cry, a shrill scream and lasted for hours and nothing would help."
<b>Collapse/Shock</b>	"She turned white with a blue tinge around her mouth and went completely limp."
<b>Excessive Sleepiness</b>	"He passed out and we couldn't wake him to feed or do anything for over 12 hours."
<b>Convulsion</b>	"Her eyes twitched, her chin trembled, her body went rigid and then would shake."
<b>Brain Inflammation</b>	"He just laid in his crib with his eyes wide open then would arch his back and scream and go un conscious. Now he has seizures."
<b>Behavior Changes</b>	"She won't sleep or eat. She throws herself down and screams for no reason. She was sweet and happy and is now out of control. She changed into a totally different child."
<b>Mental/Physical Regression</b>	"My 18 month old son stopped talking and walking after those shots. He developed severe allergies, constant diarrhea, ear infections and was sick all the time."
<b>Other reported vaccine reactions include:</b> loss of muscle control, paralysis, regressive autism, asthma, arthritis, blood disorders, diabetes, Guillain Barre syndrome, sudden death.	
<i>*Call a doctor immediately or go to an emergency room if symptoms of serious vaccine reaction complications or dramatic changes in physical, mental, or emotional behavior occur after vaccination.</i>	

**NATIONAL CHILDHOOD VACCINE INJURY ACT OF 1986**

Under the federal vaccine injury compensation program (VICP), more than \$2.5 billion has been paid to vaccine injured individuals, as well as to families, whose children have died after vaccination in the U.S.

**REPORT VACCINE REACTIONS**

Serious vaccine reactions should be documented in medical records and promptly reported to the federal Vaccine Adverse Events Reporting System (VAERS).

**LEARN MORE**

Go to [NVIC.org](http://NVIC.org) to learn more about signs and symptoms of infectious diseases and vaccine reactions; how to report vaccine reactions; how to apply for federal vaccine injury compensation; and how to protect your legal right to informed consent to vaccination in America.

# Vaccination.

Your health. Your family. Your choice.



**Is the Childhood Vaccine Schedule Safe?**

An epidemic of chronic disease and disability is plaguing America's children, who are the most highly vaccinated children in the world and also among the most chronically ill and disabled. Today, the Centers for Disease Control (CDC) states that 1 child in 6 in America suffers with learning disabilities while millions more suffer with asthma, diabetes and other chronic allergic and autoimmune diseases. The epidemic of chronic disease and disability among children has increased dramatically in the past five decades.

**U.S. CHILD CHRONIC DISEASE INCREASES**

<b>1976:</b> 1 child in 30 was learning disabled	→	<b>2013:</b> 1 child in 6 is learning disabled.
<b>1980:</b> 1 child in 27 had asthma	→	<b>2013:</b> 1 child in 9 has asthma.
<b>1990's:</b> 1 child in 555 developed autism	→	<b>2013:</b> 1 child in 50 develops autism.
<b>2001:</b> 1 child in 500 had diabetes	→	<b>2013:</b> 1 child in 400 has diabetes.

**THREE TIMES AS MANY VACCINATIONS FOR CHILDREN**

- 1953:** CDC recommended 16 doses of 4 vaccines (smallpox, DPT) between two months and age six.
- 1983:** CDC recommended 23 doses of 7 vaccines (DPT, MMR, polio) between two months and age six.
- 2013:** CDC recommended 49 doses of 14 vaccines between day of birth and age six and 69 doses of 16 vaccines between day of birth and age 18.

**MULTIPLE VACCINATIONS GIVEN SIMULTANEOUSLY**

In 1983, the CDC directed doctors to give a child no more than 4 vaccines (DPT, polio) simultaneously. By 2013, the CDC directed that a child can receive 8 or more vaccines at once.

The Institute of Medicine published a report in 2013 stating that "key elements of the entire [child vaccine] schedule – the number, frequency, timing, order and age of administration of vaccines – have not been systematically examined in research studies."

**VACCINATIONS DURING PREGNANCY**

A new CDC policy directs doctors to give pregnant women one dose of influenza vaccine in any trimester and one dose of pertussis containing Tdap vaccine after 20 weeks during every pregnancy. The U.S. Food and Drug Administration (FDA) has determined that there are no adequate, well controlled studies conducted in pregnant women to determine if it is safe for the developing fetus or pregnant woman to receive Tdap and Influenza vaccines during pregnancy.

