

# Immunization Requirements for Childcare/Preschool Children

<b>Age 3 months</b>	<b>Age 5 months</b>	<b>Age 7 months</b>	<b>Age 12 months</b>	<b>Age 16 months</b>	<b>Age 19 months</b>	<b>Age 4 years</b>	<b>Age 5 years</b>
<u>DTaP</u> 1 dose	<u>DTaP</u> 2 doses	<u>DTaP</u> 3 doses	<u>DTaP</u> 3 doses	<u>DTaP</u> 4 doses	<u>DTaP</u> 4 doses	* <u>DTaP</u> 4 or 5 doses	* <u>DTaP</u> 4 or 5 doses
<u>Polio (IPV)</u> 1 dose	<u>Polio (IPV)</u> 2 doses	<u>Polio (IPV)</u> 2 doses	<u>Polio (IPV)</u> 2 doses	<u>Polio (IPV)</u> 2 doses	<u>Polio (IPV)</u> 3 doses	* <u>Polio (IPV)</u> 3 or 4 doses	* <u>Polio (IPV)</u> 3 or 4 doses
<u>Hib</u> 1 dose	<u>Hib</u> 2 dose	<u>Hib</u> 2 doses	* <u>Hib</u> 3 doses	* <u>Hib</u> 3 or 4 doses	* <u>Hib</u> 3 or 4 doses	* <u>Hib</u> 3 or 4 doses	* <u>Hib</u> – no requirement
<u>Hepatitis B</u> 1 dose	<u>Hepatitis B</u> 2 doses	<u>Hepatitis B</u> 2 doses	<u>Hepatitis B</u> 2 doses	<u>Hepatitis B</u> 2 doses	<u>Hepatitis B</u> 3 doses	<u>Hepatitis B</u> 3 doses	<u>Hepatitis B</u> 3 doses
<u>Pneumococcal (PCV)</u> 1 dose	<u>Pneumococcal (PCV)</u> 2 doses	* <u>Pneumococcal (PCV)</u> 3 doses	* <u>Pneumococcal (PCV)</u> 4 doses	* <u>Pneumococcal (PCV)</u> 4 doses	* <u>Pneumococcal (PCV)</u> 4 doses	* <u>Pneumococcal (PCV)</u> 4 doses	* <u>Pneumococcal (PCV)</u> – no requirement
			<u>Hepatitis A</u> 1 dose effective for the school year beginning on or after <b>July 1, 2018</b>	<u>Hepatitis A</u> 1 dose effective for the school year beginning on or after <b>July 1, 2018</b>	* <u>Hepatitis A</u> 2 doses effective for the school year beginning on or after <b>July 1, 2018</b>	* <u>Hepatitis A</u> 2 doses effective for the school year beginning on or after <b>July 1, 2018</b>	* <u>Hepatitis A</u> 2 doses effective for the school year beginning on or after <b>July 1, 2018</b>
				<u>MMR</u> 1 dose	<u>MMR</u> 1 dose	<u>MMR</u> 2 doses	<u>MMR</u> 2 doses
				<u>Varicella</u> 1 dose	<u>Varicella</u> 1 dose	<u>Varicella</u> 2 doses	<u>Varicella</u> 2 doses

\*DTaP - Fifth dose not needed if dose #4 given after 4<sup>th</sup> B-day and 6 months from previous dose

\*Polio - Fourth dose not needed if dose #3 given after 4<sup>th</sup> B-day and 6 months from previous dose

\*Hib - last dose must be on or after 12 months of age-not required at age 5 years and older; number of total doses (3 or 4) depends on brand used

\*PCV - last dose must be on or after 12 months of age-not required at age 5 years and older

\*Varicella - no doses needed if history of Chickenpox or Zoster disease is verified by a medical provider

\*Hepatitis A – 2 doses spaced 6 months apart- effective for the school year beginning on or after **July 1, 2018**