Multiple Vaccinations Given Simultaneously

In 1983, the CDC directed doctors to give a child no more than 4 vaccines (DPT, Polio) simultaneously. By 2013, the CDC directed that a child can receive 8 or more vaccines at once.

The Institute of Medicine published a report in 2013 stating that "key elements of the entire (CDC recommended childhood vaccine) schedule - the number, frequency, timing, order and age of administration of vaccines - have not been systematically examined in research studies."

Vaccinations During Pregnancy

A new CDC policy directs doctors to give pregnant women one dose of influenza vaccine in any trimester and one dose of pertussis containing Tdap vaccine after 20 weeks during every pregnancy.

The Food and Drug Administration (FDA) has determined that large, well controlled long-term studies have not been conducted to confirm that influenza and Tdap vaccination during pregnancy is safe.

ABOUT US

The non-profit National Vaccine Information Center (NVIC) is an independent clearinghouse for information on diseases and vaccine science, policy, law and the ethical principle of informed consent. NVIC publishes information about vaccination and health to encourage educated decisionmaking. NVIC does not make vaccine use recommendations. NVIC supports the availability of all preventive health care options and the legal right for individuals to make informed, voluntary health choices for themselves and their children.

LEARN MORE

Go to NVIC.org and learn more about signs and symptoms of infectious diseases and vaccine reactions; how to report vaccine reactions; how to meet deadlines for applying for federal vaccine injury compensation and how to protect your legal right to informed consent to vaccination in America.

CONTACT US

21525 Ridgetop Circle, Suite 100 Sterling, Virginia 20166 ContactUs@nvic.org

> National Vaccine Information Center NVIC.org & NVICAdvocacy.org

Before You Take A Risk, Find Out What It Is

Based on the CDC's 2023

RECOMMENDED Childhood Vaccine Schedule





Doses of 15 Vaccines Before age 6





Doses of 17 Vaccines Before age 18

BEFORE YOU TAKE A RISK, FIND OUT WHAT IT IS

Birth (12 Hours)

Hepatitis B

Two Months

Diphtheria Tetanus Pertussis Polio Haemophilus Influenzae Type B (HIB) Rotavirus Hepatitis B Pneumococcal (PCV)

Four Months

Diphtheria Tetanus Pertussis Polio HIB Rotavirus PCV

0



Six Months

Diphtheria Tetanus Pertussis Polio HIB Rotavirus Hepatitis B PCV Influenza COVID-19

Seven Months

Influenza COVID-19

Twelve to Eighteen Months

Diphtheria Tetanus Pertussis Measles Mumps Rubella HIB PCV Varicella Hepatitis A (2)

Two to Six Years

Diphtheria Tetanus Pertussis Polio Measles Mumps Rubella Varicella Influenza (5)

Seven to Eighteen Years

Diphtheria Tetanus Pertussis Influenza (12) Human Papillovirus (HPV) (2) Meningococcal (2) COVID-19

Vaccine Ingredients: Different vaccine contain different ingredients including lab altered live or inactivated viruses and bacteria, chemicals, metals, proteins, antibiotics and human, animal and insect DNA and RNA.